

STABILITY OF A TREE-SHAPED NETWORK OF STRINGS AND BEAMS

KAÏS AMMARI AND FARHAT SHEL

ABSTRACT. In this paper we study the stability of a tree-shaped network of elastic strings and beams with some feedbacks at the ends. The whole system is asymptotically stable. Moreover, the energy of the solution decay exponentially to zero if there is no beam following a string (from the root to the leaves) and decay polynomially if not. We use a frequency domain method from the semigroups theory.

CONTENTS

1. Introduction	2
2. Abstract setting	4
3. Asymptotic behavior	7
3.1. Asymptotic stability	8
3.2. Exponential stability	12
3.3. Polynomial stability	16
Comment	20
Acknowledgements	21
References	21

2010 *Mathematics Subject Classification.* 35R02, 35M33, 35B40, 93D15, 74K05, 74K10.

Key words and phrases. uniform stability, polynomial stability , frequency domain method, network, strings, Euler-Bernoulli beams.

1. INTRODUCTION

In past decades, the dynamic behavior of networks of flexible structures have been studied by some authors. See for instance, [18, 13, 9] and the references therein. The importance of these studies lies in the need for engineering to eliminate vibrations in such composite structures. In this paper we consider a model ($\mathcal{S}$) of a tree-shaped network of N elastic materials, constituting of **strings** and Euler-Bernoulli **beams**.

To start let us first introduce some notations for the tree under consideration (as introduced in [1] or in [21]). Let $\mathcal{G}$ be a finite planar tree, with N edges and $p = N + 1$ vertices. We denote by $E = \{e_1, \dots, e_N\}$ the set of edges, $\mathcal{V} = \{a_1, \dots, a_p\}$, the set of vertices of $\mathcal{G}$ and we suppose that a_1 is the root of $\mathcal{G}$ and that a_1 and a_2 are ends of e_1 .

For a fixed vertex a_k , let $I(a_k)$ be the set of indices of edges adjacent to a_k . Denote by $\mathcal{V}_{int}$ the set of interior vertices and $\mathcal{V}_{ext}$ the set of exterior vertices of $\mathcal{G}$. Each edge e_j of length ℓ_j is a curve parametrized by

$$\pi_j : [0, \ell_j] \longrightarrow e_j, \quad x_j \mapsto \pi_j(x_j),$$

and sometimes identified with the interval $(0, \ell_j)$. The incidence matrix $D = (d_{ij})_{p \times N}$ is defined by

$$d_{ij} = \begin{cases} 1 & \text{if } \pi_j(\ell_j) = a_i, \\ -1 & \text{if } \pi_j(0) = a_i, \\ 0 & \text{otherwise,} \end{cases}$$

and for a function $u : \mathcal{G} \longrightarrow \mathbb{C}$ we set $u^j = u \circ \pi_j$ its restriction to the edge e_j and we will denote $u^j(x) = u^j(\pi_j(x))$ for any x in $(0, \ell_j)$.

We denote by $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ the inner product and norm in L^2 -space, respectively.

Suppose that the equilibrium position of the tree of elastic strings and beams coincides with $\mathcal{G}$.

Our model is then described as follows: every string e_j satisfies the following equation

$$(1) \quad u_{tt}^j - u_{xx}^j = 0 \text{ in } (0, \ell_j) \times (0, \infty),$$

and every beam e_j satisfies the following equation

$$(2) \quad u_{tt}^j + u_{xxxx}^j = 0 \text{ in } (0, \ell_j) \times (0, \infty)$$

where $u^j = u^j(x, t)$ is the function describing the displacement of the string or beam e_j .

The initial conditions are

$$(3) \quad u^j(x, 0) = u_0^j(x), \quad u_t^j(x, 0) = u_1^j(x).$$

Denote by $I_S(a_k)$ (resp. $I_B(a_k)$) the set of strings (resp. beams) adjacent to a_k and by $\mathcal{V}_{ext}^S$ and $\mathcal{V}_{ext}^B$ respectively the set of external nodes of strings and those of beams, different from a_1 . Then the transmission conditions at the inner nodes are

$$(4) \quad \begin{cases} u^j(a_k, t) = u^l(a_k, t), & j, l \in I(a_k), \quad a_k \in \mathcal{V}_{int}, \\ u_{xx}^j(a_k, t) = u_{xx}^l(a_k, t), & j, l \in I_B(a_k), \quad a_k \in \mathcal{V}_{int}, \\ \sum_{j \in I_B(a_k)} d_{kj} u_x^j(a_k, t) = 0, & a_k \in \mathcal{V}_{int}, \\ \sum_{j \in I_B(a_k)} d_{kj} u_{xxx}^j(0, t) - \sum_{j \in I_S(a_k)} d_{kj} u_x^j(0, t) = 0, & a_k \in \mathcal{V}_{int}, \end{cases}$$

and the boundary conditions are

$$(5) \quad \begin{cases} u^{j_k}(a_k, t) = 0, & a_k \in \mathcal{V}_{ext}, \\ u_{xx}^{j_k}(a_k, t) = 0, & a_k \in \mathcal{V}_{ext}^B \end{cases}$$

where j_k is the index of the unique edge adjacent to $a_k \in \mathcal{V}_{ext}$.

For a classical solution u of $(\mathcal{S})$, the energy is defined as the sum of the energy of its components, that is,

$$\begin{aligned} E(t) &= \frac{1}{2} \sum_{j=1}^N \int_0^{\ell_j} |u_t^j(x, t)|^2 dx + \frac{1}{2} \sum_{j \in I_S} \int_0^{\ell_j} |u_x^j(x, t)|^2 dx \\ &\quad + \frac{1}{2} \sum_{j \in I_B} \int_0^{\ell_j} |u_{xx}^j(x, t)|^2 dx \end{aligned}$$

where I_S and I_B are the sets of indices of strings and beams respectively. Differentiate formally the energy function with respect to time t , we get

$$\frac{dE}{dt}(t) = 0,$$

and the system is conservative.

Stability of such models of networks of strings or of beams, has been proved before, by applying a control at an external node or by forcing the damping conditions at inner nodes. In [4], the authors proved the polynomial stability of a star-shaped network of strings when a feedback is applied at the common node and in [5] and [26] the authors proved a similar result for a tree of strings when the feedback is applied at an external node. In [2] we consider a network of beams. See also [27] for exponential stability of a star-shaped network of beams and [16] for asymptotic stability of a star-shaped network of Timoshenko beams. In [24] and [25] we add thermoelastic edges to the network of elastic materials to obtain an exponential stability result.

For strings-beams networks see [6] where the authors considered a star-shaped network of beams and a string, with controls applied at all the exterior nodes. They proved a result of exponential stability. Some results of polynomial stability have proved before, for coupled string-beam systems [3] (see [9] for general setting) and for chains of alternated beams and strings [7], when feedbacks are applied at inner nodes. The case of a $2-d$ coupled system of a wave equation and a plate equation has been studied by K. Ammari and S. Nicaise in [8]. They proved a result of exponential stability under some geometric conditions.

In this paper we study a more general case of networks, in fact, it is the model $(\mathcal{S})$ presented at the beginning, stabilized by applying feedbacks at all leaves (the root remains free). For this, let $\mathcal{V}^* = \mathcal{V} - \{a_1\}$, $\mathcal{V}_{ext}^* = \mathcal{V}_{ext} - \{a_1\}$ and let δ in $\{0, 1\}$ with $\delta = 1$ if e_1 is a string and $\delta = 0$ if e_1 is a beam. Then instead of (5) we take

$$\begin{aligned} u^1(a_1, t) &= 0, \quad (1 - \delta)u_{xx}^1(a_1, t) = 0, \\ u_x^{j_k}(a_k, t) &= -d_{kj}u_t^{j_k}(a_k, t), \quad a_k \in \mathcal{V}_{ext}^S, \\ u_{xxx}^{j_k}(a_k, t) &= d_{kj}u_t^{j_k}(a_k, t), \quad u_x^{j_k}(a_k, t) = 0, \quad a_k \in \mathcal{V}_{ext}^B. \end{aligned}$$

Formally, we have

$$\frac{d}{dt}E(t) = - \sum_{a_k \in \mathcal{V}_{ext}^*} |u_t^{j_k}(a_k, t)|^2 \leq 0.$$

So the system is dissipative.

We prove different decay results of the energy of the system depending on the position of beams relative to strings. Precisely $(\mathcal{S})$ is exponentially stable if there is no beam following a string from the root to leaves and polynomially stable if not. Moreover, we give an example corresponding to the last case which is not exponentially stable. The method that we use to show exponential or polynomial stability is based on the resolvent approach.

The paper is organized as follows: In section 2, we reformulate the system $(\mathcal{S})$ as an evolution equation in a Hilbert space, and prove that it is associated with a $\mathcal{C}_0$ -semigroup of contraction and in section 3, by using frequency domain method, we first prove, under some conditions, that the system $(\mathcal{S})$ is exponentially stable, then we give a result of polynomial stability.

2. ABSTRACT SETTING

The aim of this section is to rewrite the system $(\mathcal{S})$ as an evolution equation in an appropriate Hilbert space. We then prove the existence and uniqueness of solutions of the problem using semigroup theory.

Let us consider

$$V = \left\{ f = (f_S, f_B) \in \prod_{j \in I_S} H^1(0, \ell_j) \times \prod_{j \in I_B} H^2(0, \ell_j) \mid f \text{ satisfies (6)} \right\}$$

where $f_S = (f_j)_{j \in I_S}$, $f_B = (f_j)_{j \in I_B}$, and

$$(6) \quad \begin{cases} f^1(a_1) = 0, \\ f^j(a_k) = f^l(a_k), \quad j, l \in I(a_k), \quad a_k \in \mathcal{V}_{int}, \\ \partial_x f^{j_k}(a_k) = 0, \quad a_k \in \mathcal{V}_{ext}^B, \\ \sum_{j \in I_B(a_k)} d_{kj} \partial_x f^j(a_k) = 0, \quad a_k \in \mathcal{V}_{int}. \end{cases}$$

Note that we can rewrite the last two equations in one, as follows

$$\sum_{j \in I_B(a_k)} d_{kj} \partial_x f^j(a_k) = 0, \quad a_k \in \mathcal{V}^*.$$

Define the energy space of $(\mathcal{S})$ by

$$\mathcal{H} = V \times \prod_{j=1}^N L^2(0, \ell_j)$$

endowed by the inner product

$$\langle y_1, y_2 \rangle_{\mathcal{H}} := \sum_{j \in I_S} \langle \partial_x f_1^j, \partial_x f_2^j \rangle + \sum_{j \in I_B} \langle \partial_x^2 f_1^j, \partial_x^2 f_2^j \rangle + \sum_{j=1}^N \langle g_1^j, g_2^j \rangle$$

where $y_k = (f_k, g_k)$, $k = 1, 2$. Then $\mathcal{H}$ is a Hilbert space.

Now define the operator $\mathcal{A}$ on $\mathcal{H}$ by

$$\mathcal{D}(\mathcal{A}) = \left\{ y = (u, v) \in V \times V \mid u_S \in \prod_{j \in I_S} H^2(0, \ell_j), u_B \in \prod_{j \in I_B} H^4(0, \ell_j) \text{ and } y \text{ satisfies (7)} \right\}$$

where

$$(7) \quad \begin{cases} \partial_x u^{j_k}(a_k) = -d_{kj} v^{j_k}(a_k), & a_k \in \mathcal{V}_{ext}^S, \\ (1 - \delta) \partial_x^2 u^1(a_1) = 0, \\ \partial_x^2 u^j(a_k) = \partial_x^2 u^l(a_k), & j, l = I_B(a_k), \quad a_k \in \mathcal{V}_{int}, \\ \partial_x^3 u^{j_k}(a_k) = d_{kj} v^{j_k}(a_k), & a_k \in \mathcal{V}_{ext}^B, \\ \sum_{j \in I_S(a_k)} d_{kj} \partial_x u^j(a_k) - \sum_{j \in I_B(a_k)} d_{kj} \partial_x^3 u^j(a_k) = 0, & a_k \in \mathcal{V}_{int} \end{cases}$$

and

$$\mathcal{A} \begin{pmatrix} u_S \\ u_B \\ v_S \\ v_B \end{pmatrix} = \begin{pmatrix} v_S \\ v_B \\ \partial_x^2 u_S \\ -\partial_x^4 u_B \end{pmatrix}, \quad (u_S, u_B, v_S, v_B) \in (\mathcal{A}).$$

Then, putting $y = (u, u_t)$, we write the system (S) into the following first order evolution equation

$$(8) \quad \begin{cases} \frac{dy}{dt} = \mathcal{A}y, \\ y(0) = y_0 \end{cases}$$

on the energy space $\mathcal{H}$, where $y_0 = (u_0, u_1)$.

We have the following result,

Lemma 1. *The operator $\mathcal{A}$ is the infinitesimal generator of a $\mathcal{C}_0$ -semigroup of contraction $(T(t))_{t \geq 0}$.*

Proof. By Lumer-Phillips' theorem (see [22]), it suffices to show that $\mathcal{A}$ is dissipative maximal. First, for any $y \in \mathcal{H}$ we have,

$$\begin{aligned} \operatorname{Re}(\langle \mathcal{A}y, y \rangle_{\mathcal{H}}) &= \operatorname{Re} \left(\sum_{j \in I_S} \int_0^{\ell_j} (\partial_x v^j \overline{\partial_x u^j} dx + \partial_x^2 u^j \overline{v^j}) dx \right. \\ &\quad \left. + \sum_{j \in I_B} \int_0^{\ell_j} (\partial_x^2 v^j \overline{\partial_x^2 u^j} dx - \partial_x^4 u^j \overline{v^j}) dx \right). \end{aligned}$$

By some integrations by parts, we obtain using the boundary and transmissions conditions (6-7),

$$\operatorname{Re}(\langle \mathcal{A}y, y \rangle_{\mathcal{H}}) = - \sum_{a_k \in \mathcal{V}_{ext}^*} |v^{j_k}(\ell_{j_k})|^2 \leq 0.$$

Then the operator $\mathcal{A}$ is dissipative.

We show now that every positive real number λ belongs to $\rho(\mathcal{A})$. Let $z = (f, g) \in \mathcal{H}$, we look for $y = (u, v) \in \mathcal{D}(\mathcal{A})$ such that

$$(\lambda - \mathcal{A})y = z$$

i.e.,

$$(9) \quad \lambda u^j - v^j = f^j, \quad j = 1, \dots, N,$$

$$(10) \quad \lambda v^j - \partial_x^2 u^j = g^j, \quad j \in I_S,$$

$$(11) \quad \lambda v^j + \partial_x^4 u^j = g^j, \quad j \in I_B.$$

Then

$$(12) \quad \lambda^2 u^j - \partial_x^2 u^j = g^j + \lambda f^j, \quad j \in I_S,$$

$$(13) \quad \lambda^2 u^j + \partial_x^4 u^j = g^j + \lambda f^j, \quad j \in I_B.$$

Let w in V . Multiplying the first equation by w_S and the second equation by w_B , and integrating by parts, we get respectively

$$\lambda^2 \int_0^{\ell_j} u^j \overline{w^j} dx + \int_0^{\ell_j} \partial_x u^j \partial_x \overline{w^j} dx - \partial_x u^j \overline{w^j} \Big|_0^{\ell_j} = \int_0^{\ell_j} (g^j + \lambda f^j) \overline{w^j} dx$$

for j in I_S and

$$\lambda^2 \int_0^{\ell_j} u^j \overline{w^j} dx + \int_0^{\ell_j} \partial_x^2 u^j \partial_x^2 \overline{w^j} dx + \partial_x^3 u^j \overline{w^j} \Big|_0^{\ell_j} - \partial_x^2 u^j \partial_x \overline{w^j} \Big|_0^{\ell_j} = \int_0^{\ell_j} (g^j + \lambda f^j) \overline{w^j} dx$$

for j in I_B . Now summing the two obtained equations, the left hand side will be

$$\begin{aligned} & \lambda^2 \sum_{j=1}^N \int_0^{\ell_j} u^j \overline{w^j} dx + \sum_{j \in I_S} \int_0^{\ell_j} \partial_x u^j \partial_x \overline{w^j} dx + \sum_{j \in I_B} \int_0^{\ell_j} \partial_x^2 u^j \partial_x^2 \overline{w^j} dx \\ & + \sum_{a_k \in \mathcal{V}_{int}} \left(- \sum_{j \in I_S(a_k)} d_{kj} \overline{w^j}(a_k) \partial_x u^j(a_k) + \sum_{j \in I_B(a_k)} d_{kj} \overline{w^j}(a_k) \partial_x^3 u^j(a_k) \right) \\ & + \sum_{a_k \in \mathcal{V}_{ext}} v^{jk}(a_k) \overline{w^{jk}}(a_k) + \sum_{a_k \in \mathcal{V}_{int}} \sum_{j \in I_B(a_k)} d_{kj} \partial_x^2 u^j(a_k) \partial_x \overline{w^j}(a_k). \end{aligned}$$

We find, by taking into account (6) and (7)

$$(14) \quad a(u, w) = F(w)$$

where

$$\begin{aligned} a(u, w) &= \lambda^2 \sum_{j=1}^N \int_0^{\ell_j} u^j \overline{w^j} dx + \sum_{j \in I_S} \int_0^{\ell_j} \partial_x u^j \partial_x \overline{w^j} dx + \sum_{j \in I_B} \int_0^{\ell_j} \partial_x^2 u^j \partial_x^2 \overline{w^j} dx \\ &+ \lambda \sum_{a_k \in \mathcal{V}_{ext}} u^{jk}(a_k) \overline{w^{jk}}(a_k) \end{aligned}$$

and

$$F(w) = \sum_{j=1}^N \int_0^{\ell_j} (g^j + \lambda f^j) \overline{w^j} dx + \lambda \sum_{a_k \in \mathcal{V}_{ext}} f^{jk}(a_k) \overline{w^{jk}}(a_k).$$

a is a continuous sesquilinear form on $V \times V$ and F is a continuous anti-linear form on V . Moreover, there exists $C > 0$ such that, for every $w \in V$

$$|a(w, w)| \geq C \|w\|_V^2,$$

where

$$\|u\|_V^2 = \sum_{j=1}^N \int_0^{\ell_j} |u^j|^2 dx + \sum_{j \in I_S} \int_0^{\ell_j} |\partial_x u^j|^2 dx + \sum_{j \in I_B} \int_0^{\ell_j} |\partial_x^2 u^j|^2 dx.$$

By the Lax-Milgram's lemma, problem (14) has a unique solution u in V . It is easy to verify that: u belongs to $\prod_{j \in I_S} H^2(0, \ell_j) \times \prod_{j \in I_B} H^4(0, \ell_j)$, $v = \lambda u - f \in V$, u_S and




FIGURE 1. first tree




FIGURE 2. second tree

u_B satisfies respectively (10) and (11), and the conditions

$$\begin{cases} \partial_x u^{jk}(a_k) = -d_{kj} v^{jk}(a_k), & a_k \in \mathcal{V}_{ext}^S, \\ (1 - \delta) \partial_x^2 u^1(\ell_1) = 0, \\ \partial_x^2 u^j(a_k) = \partial_x^2 u^l(a_k), & j, l = I_B(a_k), \quad a_k \in \mathcal{V}_{int}, \\ \partial_x^3 u^{jk}(a_k) = d_{kj} v^{jk}(a_k), & a_k \in \mathcal{V}_{ext}^B, \\ \sum_{j \in I_S(a_k)} d_{kj} \partial_x u^j(a_k) - \sum_{j \in I_B(a_k)} d_{kj} \partial_x^3 u^j(a_k) = 0, & a_k \in \mathcal{V}_{int}. \end{cases}$$

Furthermore

$$\|y\|_{\mathcal{H}}^2 \leq c \|z\|_{\mathcal{H}}^2,$$

where c is a positive constant independent of y . In conclusion, $y = (u, v) \in \mathcal{D}(\mathcal{A})$ and $(\lambda - \mathcal{A})^{-1} \in \mathcal{L}(\mathcal{H})$, that is, $\lambda \in \rho(\mathcal{A})$. $\square$

Corollary 2. *For an initial datum $y_0 \in \mathcal{H}$ there exists a unique solution $y \in C([0, +\infty), \mathcal{H})$ of problem (8). Moreover if $y_0 \in \mathcal{D}(\mathcal{A})$, then $y \in C([0, +\infty), \mathcal{D}(\mathcal{A})) \cap C^1([0, +\infty), \mathcal{H})$.*

Remark 3. *Note that, by the Sobolev embedding theorem, $(I - \mathcal{A})^{-1}$ is a compact operator and then the spectrum of $\mathcal{A}$ consists of all isolated eigenvalues, i.e., $\sigma(\mathcal{A}) = \sigma_p(\mathcal{A})$.*

3. ASYMPTOTIC BEHAVIOR

The aim of this section is to show that the system $(\mathcal{S})$ is asymptotically stable. Moreover, we will prove that the solution is exponentially stable if there is no beam following a string, from the root to the leaves, as in the first tree (Figure 1), and polynomially stable if at least a beam follows a string, as in the second tree (Figure 2). Finally, The lack of exponential stability is proved on an example.

We will use the following two results. The first gives us a necessary and sufficient condition for the exponential stability of a $\mathcal{C}_0$ -semigroup of contraction, for the proof see [15], [17] or [23].

Theorem 4. A $\mathcal{C}_0$ -semigroup of contraction $e^{t\mathcal{L}}$ on a Hilbert space is exponentially stable if, and only if,

$$(15) \quad i\mathbb{R} \cap \sigma(\mathcal{L}) = \emptyset$$

and

$$(16) \quad \lim_{|\beta| \rightarrow \infty} \sup \| (i\beta - \mathcal{L})^{-1} \| < \infty.$$

The second, due to [12] (see also [10] and [19]), characterizes the polynomial decay of a $\mathcal{C}_0$ -semigroup of contraction.

Theorem 5. A $\mathcal{C}_0$ -semigroup of contraction $e^{t\mathcal{L}}$ on a Hilbert space $\mathcal{H}$ satisfies

$$\|e^{t\mathcal{L}}y_0\| \leq \frac{C}{t^\alpha} \|y_0\|_{\mathcal{D}(\mathcal{L})}$$

for some constant $C > 0$ and for $\alpha > 0$ if, and only if, (15) holds and

$$(17) \quad \lim_{|\beta| \rightarrow \infty} \sup \frac{1}{\beta^\alpha} \| (i\beta - \mathcal{L})^{-1} \| < \infty.$$

3.1. Asymptotic stability. In this section, we prove (15), that is, the system $(\mathcal{S})$ is asymptotically stable.

Theorem 6. The semigroup $(T(t))_{t \geq 0}$, generated by the operator $\mathcal{A}$ is asymptotically stable.

Proof. It suffices to show that (15) holds, otherwise, by taking into account Remark 3, there is a real number β , such that $\lambda := i\beta$ is an eigenvalue of $\mathcal{A}$. Let $y = (u, v)$ the corresponding eigenvector. We have

$$(18) \quad \begin{cases} v^j = \lambda u^j & \text{for } j \text{ in } \{1, \dots, N\}, \\ \partial_x^2 u^j = \lambda v^j & \text{for } j \text{ in } I_S, \\ -\partial_x^4 u^j = \lambda v^j & \text{for } j \text{ in } I_B. \end{cases}$$

If $\lambda = 0$, multiplying the second and the third equations of (18) by u^j and summing, we obtain, using (6) and (7),

$$\sum_{j \in I_S} \|\partial_x u^j\|^2 + \sum_{j \in I_B} \|\partial_x^2 u^j\|^2 = 0$$

which implies that $u = 0$ and hence $v = 0$. Then, in the sequel, we suppose that $\lambda \neq 0$.

Taking the real part of the inner product of $\lambda y - \mathcal{A}y = 0$ with y in $\mathcal{H}$, we obtain

$$\operatorname{Re}(\langle \mathcal{A}y, y \rangle_{\mathcal{H}}) = - \sum_{a_k \in \mathcal{V}_{ext}^*} |v^{j_k}(a_k)|^2 = 0.$$

Thus $v^{j_k}(a_k) = 0$ for $a_k \in \mathcal{V}_{ext}^*$ and then $u^{j_k}(a_k) = 0$ for $a_k \in \mathcal{V}_{ext}^*$, which holds for $k = 1$; $\partial_x^3 u^{j_k}(a_k) = 0$ for $a_k \in \mathcal{V}_{ext}^B$ and $\partial_x u^{j_k}(a_k) = 0$ for $a_k \in \mathcal{V}_{ext}^S$.

Then, u is zero on every edge attached to a leaf, and by iteration, on every maximal subgraph of strings not followed by beams.

Now let $\mathcal{G}'$ a maximal subgraph of beams not followed by strings. We want to prove that u is zero on $\mathcal{G}'$.

First case: $\mathcal{G}' = \mathcal{G}$. For each j in $\{1, \dots, N\}$, substituting the first equation of (18) into the third, we obtain,

$$(19) \quad \partial_x^4 u^j + \lambda^2 u^j = 0.$$

For the sequel, we use a matrix method [11]. First of all we shall introduce some definitions and notations used in [11] (see also [1]).

The adjacency matrix $E = (e_{jk})_{p \times p}$ of $\mathcal{G}$ is defined by

$$e_{jk} = \begin{cases} 1 & \text{if } a_j \text{ and } a_k \text{ are adjacent,} \\ 0 & \text{otherwise.} \end{cases}$$

The Hadamard product of two matrices $A = (a_{jk})_{p \times p}$ and $B = (b_{jk})_{p \times p}$ is the matrix $A * B = (a_{jk}b_{jk})_{p \times p}$. For a function $r : \mathbb{R} \rightarrow \mathbb{R}$, we define the matrix $r(A) = (r_{jk})_{p \times p}$ by

$$r_{jk} = \begin{cases} r(a_{jk}) & \text{if } e_{jk} = 1, \\ 0 & \text{otherwise,} \end{cases}$$

in particular, if $r(x) = x^q$ then, we write $A^{(q)}$ instead of $r(A)$. Furthermore, the matrix $L = (\ell_{jk})_{p \times p}$ is defined as follows

$$\ell_{jk} = \begin{cases} \ell_{s(j,k)} & \text{if } e_{jk} = 1, \\ 0 & \text{otherwise,} \end{cases}$$

where $s(j, k)$ is the indice of the edge connecting a_j to a_k .

To a function f on $\mathcal{G}$ is associated the matrix function F defined by

$$F : [0, 1] \longrightarrow \mathbb{C}^{p \times p}, x \longmapsto F(x) = (f_{jk}(x))_{p \times p},$$

with

$$f_{jk}(x) = e_{jk} f_{s(j,k)} \left[\ell_{s(j,k)} \left(\frac{1 + d_{js(j,k)}}{2} - x d_{js(j,k)} \right) \right].$$

The system (19) is then rewritten as

$$(20) \quad L^{(-4)} * U'''' + \lambda U = 0.$$

Integrate equation (20) we obtain

$$(21) \quad U(x) = A_1 * \cos(\sqrt{\beta} Lx) + A_2 * \sin(\sqrt{\beta} Lx) + B_1 * \cosh(\sqrt{\beta} Lx) + B_2 * \sinh(\sqrt{\beta} Lx),$$

where without loss of generality we have supposed that $\beta > 0$ and with $A_1, A_2, B_1, B_2 \in \mathbb{C}^{p \times p}$. Then

$$(22) \quad \begin{aligned} L^{(-1)} * U' &= \sqrt{\beta} \left(-A_1 * \sin(\sqrt{\beta} Lx) + A_2 * \cos(\sqrt{\beta} Lx) \right. \\ &\quad \left. + B_1 * \sinh(\sqrt{\beta} Lx) + B_2 * \cosh(\sqrt{\beta} Lx) \right), \end{aligned}$$

$$(23) \quad \begin{aligned} L^{(-2)} * U'' &= \beta \left(-A_1 * \cos(\sqrt{\beta} Lx) - A_2 * \sin(\sqrt{\beta} Lx) \right. \\ &\quad \left. + B_1 * \cosh(\sqrt{\beta} Lx) + B_2 * \sinh(\sqrt{\beta} Lx) \right), \end{aligned}$$

$$(24) \quad \begin{aligned} L^{(-3)} * U''' &= \beta^{3/2} \left(A_1 * \sin(\sqrt{\beta} Lx) - A_2 * \cos(\sqrt{\beta} Lx) \right. \\ &\quad \left. + B_1 * \sinh(\sqrt{\beta} Lx) + B_2 * \cosh(\sqrt{\beta} Lx) \right). \end{aligned}$$

The function U satisfies also,

$$(25) \quad U(1 - x) = U(x)^T.$$

The boundary and transmission conditions can be expressed as follows:

For the continuity condition of u at the inner nodes, there exists $\varphi = \begin{pmatrix} \varphi_1 \\ \vdots \\ \varphi_p \end{pmatrix} \in \mathbb{C}^p$

such that

$$(26) \quad U(0) = (\varphi e^T) * E,$$

where $e = \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \in \mathbb{R}^p$.

Since u is zero at all external nodes then $\varphi_j = 0$ when a_j is an external node.

The continuity condition of $\partial_x^2 u$ at the interior nodes and the fact that $\partial_x^2 u$ is zero at the root can be expressed in this manner,

there exists $\psi = \begin{pmatrix} \psi_1 \\ \vdots \\ \psi_p \end{pmatrix} \in \mathbb{C}^p$ such that $\psi_1 = 0$, and

$$(27) \quad L^{(-2)} * U''(0) = (\psi e^T) * E.$$

The forth condition of (6) and the fifth of (7) applied to u are expressed respectively as follows

$$(28) \quad (L^{(-1)} * U'(0) * E^*)e = 0$$

and

$$(29) \quad (L^{(-3)} * U'''(0) * E^*)e = 0$$

where E^* is obtained from E by annulling the first line.

Substitute (26-27) in (22-24), and (28-29) in (21-23) leads to

$$(30) \quad A_1 = \frac{1}{2}(U(0) - L^{(-2)} * U''(0)) = \frac{1}{2}((\varphi - \psi)e^T) * E,$$

$$(31) \quad B_1 = \frac{1}{2}(U(0) + L^{(-2)} * U''(0)) = \frac{1}{2}((\varphi + \psi)e^T) * E,$$

and

$$(32) \quad (A_2 * E^*)e = 0,$$

$$(33) \quad (B_2 * E^*)e = 0.$$

By taking $x = 1$ in (21) and (23) and using (25) we get, by combining the two obtained equations

$$(34) \quad \sinh^{(-1)}(\sqrt{\beta}L) * (B_1^T - B_1 \cosh(\sqrt{\beta}L)) = B_2.$$

Multiplying, in the Hadamard product, the above equation by E^* , we get, using (33)

$$(35) \quad (\sinh^{(-1)}(\sqrt{\beta}L) * (B_1^T - B_1 \cosh(\sqrt{\beta}L)) * E^*)e = 0,$$

We recall the following elementary rules for a matrix $M \in \mathbb{C}^{p \times p}$ (see [14]),

$$(36) \quad (M * B_1^T)e = M(\varphi + \psi), \quad (M * B_1)e = \text{diag}(Me)(\varphi + \psi).$$

Then, (35) implies

$$J(\varphi + \psi) = 0,$$

where

$$J = \sinh^{(-1)}(\sqrt{\beta}L) * E^* - \text{diag} \left[\left(\sinh^{(-1)}(\sqrt{\beta}L) * \cosh(\sqrt{\beta}L) * E^* \right) e \right].$$

The matrix, obtained from $J * E^{*T} * E^*$ by removing rows and columns that are zero, is a strictly diagonally dominant matrix. Since $\varphi_1 = \psi_1 = 0$, this implies that the vector $\varphi + \psi$, and hence the matrix B_1 , is zero. Return to (34) we deduce that $B_2 = 0$.

For $j = 1, \dots, N$, the expression of u^j is then

$$u^j(x) = a_1^j \cos(\sqrt{\beta}x) + a_2^j \sin(\sqrt{\beta}x), \quad (a_1^j, a_2^j \in \mathbb{C})$$

which easily implies, using the transmissions conditions and the fact that u and $\partial_x u$ vanish at the leaves, that

$$u = 0.$$

Second case: $\mathcal{G}' \neq \mathcal{G}$. Let a' be the nearest node of $\mathcal{G}'$ to a_1 . Then a' is an end of at least one string.

For simplicity of notations we will suppose, in this part, that $a' = a_1$ and $\mathcal{G}' = \mathcal{G}$ but with boundary conditions at a_1 :

$$\begin{cases} u^j(a_1) = u^l(a_1) & j, l \in I(a_1), \\ \partial_x^2 u^j(a_1) = \partial_x^2 u^l(a_1) & j, l \in I(a_1), \\ \sum_{j \in I_B(a_1)} d_{1j} \partial_x u^j(a_1) = 0. \end{cases}$$

As for the first case, there is $\psi = \begin{pmatrix} \psi_1 \\ \vdots \\ \psi_p \end{pmatrix}$ in $\mathbb{C}^p$ with $\varphi_j = 0$ when $a_j \in \mathcal{V}_{ext}^*$,

such that

$$U(0) = (\varphi e^T) * E$$

and

$$L^{(-2)} * U''(0) = (\psi e^T) * E.$$

The third and forth conditions of (6) and the fifth of (7) applied to u are expressed as follows :

$$(L^{(-1)} * U'(0) * E)e = 0$$

and

$$(L^{(-3)} * U'''(0) * E^*)e = 0.$$

As in the first case we obtain (30), (31), (32), (33) and (34). Moreover, $(\varphi_k + \psi_k)_{k=2, \dots, p}$ will be the trivial solution of a homogeneous linear system whose matrix is invertible. Then $B_1 * E^*$ and $B_2 * E^{*T} * E^*$ are zero.

For $j \in \{1, \dots, N\} - I(a_1)$ the expression of u^j is then

$$u^j(x) = a_1^j \cos(\sqrt{\beta}x) + a_2^j \sin(\sqrt{\beta}x), \quad (a_1^j, a_2^j \in \mathbb{C}),$$

which easily implies, by using the transmissions conditions and the fact that u and $\partial_x u$ vanish at the leaves,

$$u^j = 0.$$

Then we can suppose that $\mathcal{G}'$ is a start of beams e_j , $j \in I(a_1)$. Without loss of generality, we identify e_j with $(0, \ell_j)$ by taking $\pi_j(0) = a_1$. In such case we have

the following system :

$$\begin{cases} u^j(\ell_j) = 0, \quad j \in I(a_1), \\ \partial_x u^j(\ell_j) = 0 \text{ and } \partial_x^3 u^j(\ell_j) = 0, \quad j \in I(a_1), \\ u^j(0) = u^l(0) \text{ and } \partial_x^2 u^j(0) = \partial_x^2 u^l(0), \quad j, l \in I(a_1), \\ \sum_j \partial_x u^j(0) = 0. \end{cases}$$

Which implies

$$\begin{cases} a_1^j \cos(\sqrt{\beta} \ell_j) + a_2^j \sin(\sqrt{\beta} \ell_j) + b_1^j \cosh(\sqrt{\beta} \ell_j) + b_2^j \sinh(\sqrt{\beta} \ell_j) = 0, \quad j \in I(a_1), \\ -a_1^j \sin(\sqrt{\beta} \ell_j) + a_2^j \cos(\sqrt{\beta} \ell_j) = 0, \quad j \in I(a_1), \\ b_1^j \sinh(\sqrt{\beta} \ell_j) + b_2^j \cosh(\sqrt{\beta} \ell_j) = 0, \quad j \in I(a_1), \\ b_1^j = b_1^l, \quad j, l \in I(a_1), \\ \sum_j a_2^j + b_2^j = 0. \end{cases}$$

The discriminant of the above system is

$$\Delta = \sum_j \left(\prod_{k \neq j} \cosh(\sqrt{\beta} \ell_k) (\sin(\sqrt{\beta} \ell_j) + \sinh(\sqrt{\beta} \ell_j)) \right)$$

which is different from zero. We conclude that $u^j = 0$, $j \in I(a_1)$. That is, u is null on $\mathcal{G}'$.

By iteration, and using transmission conditions we conclude that u is zero on $\mathcal{G}$. The above discussion is sufficient to conclude that $y = 0$, which contradicts the fact that $y \neq 0$. $\square$

3.2. Exponential stability. In this section we suppose that there are no beam following a string (from the root to leaves), that is to say on every tree branch there is no beam between a string and a leaf (Figure 1). We prove that the solution of the whole system $(\mathcal{S})$ is exponentially stable.

Theorem 7. *If there are no beam following a string (from the root to leaves) then, the system $(\mathcal{S})$ is exponentially stable.*

Proof. It suffices to prove that (16) holds. Suppose the conclusion is false. Then there exists a sequence (β_n) of real numbers, without loss of generality, with $\beta_n \rightarrow +\infty$, and a sequence of vectors $(y_n) = (u_n, v_n)$ in $\mathcal{D}(\mathcal{A})$ with $\|y_n\|_{\mathcal{H}} = 1$, such that

$$\|(\mathbf{i}\beta_n I - \mathcal{A})y_n\|_{\mathcal{H}} \rightarrow 0$$

which is equivalent to

$$(37) \quad \mathbf{i}\beta_n u_n^j - v_n^j = f_n^j \rightarrow 0, \quad \text{in } H^1(0, \ell_j), \quad j \text{ in } I_S,$$

$$(38) \quad \mathbf{i}\beta_n u_n^j - v_n^j = f_n^j \rightarrow 0, \quad \text{in } H^2(0, \ell_j), \quad j \text{ in } I_B,$$

$$(39) \quad \mathbf{i}\beta_n v_n^j - \partial_x^2 u_n^j = g_n^j \rightarrow 0, \quad \text{in } L^2(0, \ell_j), \quad j \text{ in } I_S,$$

$$(40) \quad \mathbf{i}\beta_n v_n^j + \partial_x^4 u_n^j = g_n^j \rightarrow 0, \quad \text{in } L^2(0, \ell_j), \quad j \text{ in } I_B.$$

Then

$$(41) \quad -\beta_n^2 u_n^j - \partial_x^2 u_n^j = g_n^j + \mathbf{i}\beta_n f_n^j, \quad j \text{ in } I_S,$$

$$(42) \quad -\beta_n^2 u_n^j + \partial_x^4 u_n^j = g_n^j + \mathbf{i}\beta_n f_n^j, \quad j \text{ in } I_B$$

and

$$\|v_n^j\|^2 - \beta_n^2 \|u_n^j\|^2 \rightarrow 0, \quad j = 1, \dots, N.$$

First, Since

$$Re(\langle (\mathbf{i}\beta_n - \mathcal{A})y_n, y_n \rangle_{\mathcal{H}}) = \sum_{a_k \in \mathcal{V}_{ext}^*} |v^{j_k}(a_k)|^2,$$

we obtain

$$|v_n^{j_k}(a_k)| \longrightarrow 0, \text{ for } j \in \mathcal{V}_{ext}^*.$$

Then $\beta_n u_n^{j_k}(a_k) \longrightarrow 0$ for $a_k \in \mathcal{V}_{ext}^*$, $\partial_x^3 u_n^{j_k}(a_k) \longrightarrow 0$ for $a_k \in \mathcal{V}_{ext}^B$ and $\partial_x u_n^{j_k}(a_k) \longrightarrow 0$ for $a_k \in \mathcal{V}_{ext}^S$, and recall that $\partial_x u_n^{j_k}(a_k) = 0$ for $j \in \mathcal{V}_{ext}^B$.

Let $a_k \in \mathcal{V}_{ext}^S$ and q a function in $C^1([0, \ell_{j_k}], \mathbb{C})$. The real part of the inner product of (41) with $q \partial_x u_n^{j_k}$ gives

$$\begin{aligned} & \frac{1}{2} \beta_n^2 |u_n^{j_k}(x)|^2 q(x) \Big|_0^{\ell_{j_k}} + \frac{1}{2} |\partial_x u_n^{j_k}(x)|^2 q(x) \Big|_0^{\ell_{j_k}} \\ & - \frac{1}{2} \int_0^{\ell_{j_k}} \left(|\partial_x u_n^{j_k}(x)|^2 + \beta_n^2 |u_n^{j_k}(x)|^2 \right) \partial_x q(x) dx \\ (43) \quad & + Re \left(i \beta_n f_n^{j_k}(x) q(x) \overline{u_n^{j_k}(x)} \right) \Big|_0^{\ell_{j_k}} \longrightarrow 0. \end{aligned}$$

With $q(x) = x$ or $q(x) = \ell_{j_k} - x$ we can deduce easily that

$$(44) \quad -\frac{1}{2} \int_0^{\ell_{j_k}} \left(|\partial_x u_n^{j_k}(x)|^2 + \beta_n^2 |u_n^{j_k}(x)|^2 \right) dx \longrightarrow 0,$$

and as in [24] it follows that,

$$(45) \quad \beta_n u_n^{j_k}(a_s) \longrightarrow 0, \quad \partial_x u_n^{j_k}(a_s) \longrightarrow 0, \text{ and } Re \left(i \beta_n f_n^{j_k}(a_s) \overline{u_n^{j_k}(a_s)} \right) \longrightarrow 0$$

where a_s is the end of e_{j_k} different from a_k .

By iteration we conclude that for every $j \in I_S - \{1\}$, the properties (44) and (45) hold.

If there is no beam in the tree, then the fifth condition in (7) with (45) imply that $\partial_x u_n^1(a_2) \longrightarrow 0$. Then as for j in $\mathcal{V}_{ext}^S$, with using (45) again and the continuity condition of u at internal nodes, we obtain

$$-\frac{1}{2} \int_0^{\ell_1} \left(|\partial_x u_n^1(x)|^2 + \beta_n^2 |u_n^1(x)|^2 \right) dx \longrightarrow 0.$$

Then, we conclude that $\|y_n\| \rightarrow 0$ which contradicts the fact that $\|y_n\| = 1$ and the proof is then complete.

Now suppose that there is at least one beam. Without loss of generality, we suppose that there is no string in $\mathcal{G}$.

Let j in $\{1, \dots, N\}$ and q a function in $C^2([0, \ell_j], \mathbb{C})$ such that $\partial_x^2 q = 0$. We want to calculate the real part of the inner product of (42) by $q \partial_x u_n^j$.

Straight-forward calculations give

$$\begin{aligned} & Re \left(\langle -\beta_n^2 u_n^j, q \partial_x u_n^j \rangle \right) + Re \left(\langle \partial_x^4 u_n^j, q \partial_x u_n^j \rangle \right) = \\ & -\frac{1}{2} \beta_n^2 |u_n^j(x)|^2 q(x) \Big|_0^{\ell_j} + \frac{1}{2} \int_0^{\ell_j} \beta_n^2 |u_n^j|^2 \partial_x q dx + Re \left(\partial_x^3 u_n^j(x) q(x) \partial_x \overline{u_n^j(x)} \Big|_0^{\ell_j} \right) \\ & -\frac{1}{2} |\partial_x^2 u_n^j(x)|^2 q(x) \Big|_0^{\ell_j} + \frac{3}{2} \int_0^{\ell_j} |\partial_x^2 u_n^j|^2 \partial_x q dx - Re \left(\partial_x^2 u_n^j(x) \partial_x \overline{u_n^j(x)} \partial_x q(x) \Big|_0^{\ell_j} \right), \end{aligned}$$

and

$$\begin{aligned} \operatorname{Re} (\langle g_n^j + \mathbf{i}\beta_n f_n^j, q \partial_x u_n^j \rangle) &= \operatorname{Re} \left(\int_0^{\ell_j} g_n^j \partial_x \overline{u_n^j} q dx \right) \\ &- \operatorname{Re} \left(\mathbf{i}\beta_n \int_0^{\ell_j} \partial_x (q f_n^j) \overline{u_n^j} dx \right) + \operatorname{Re} \mathbf{i}\beta_n f_n^j(x) q(x) \overline{u_n^j}(x) \Big|_0^{\ell_j}. \end{aligned}$$

Since g_n^j , f_n^j and $\partial_x(qf_n^j)$ converge to 0 and $\mathbf{i}\beta_n u_n^j$ and $\partial_x u_n^j$ are bounded, the first and the second terms of the right member of the above equality converge to 0. It follows

$$\begin{aligned} & -\frac{1}{2}\beta_n^2 |u_n^j(x)|^2 q(x) \Big|_0^{\ell_j} + \operatorname{Re} \left(\partial_x^3 u_n^j(x) q(x) \partial_x \overline{u_n^j}(x) \Big|_0^{\ell_j} \right) - \frac{1}{2} |\partial_x^2 u_n^j(x)|^2 q(x) \Big|_0^{\ell_j} \\ & - \operatorname{Re} \left(\partial_x^2 u_n^j(x) \partial_x \overline{u_n^j} \partial_x q(x) \Big|_0^{\ell_j} \right) - \operatorname{Re} \left(\mathbf{i}\beta_n f_n^j(x) q(x) \overline{u_n^j}(x) \Big|_0^{\ell_j} \right) \\ (46) \quad & + \frac{1}{2} \int_0^{\ell_j} \beta_n^2 |u_n^j|^2 \partial_x q dx + \frac{3}{2} \int_0^{\ell_j} |\partial_x^2 u_n^j|^2 \partial_x q dx \longrightarrow 0. \end{aligned}$$

In particular if a_k is in $\mathcal{V}_{ext}^B$ then with $j = j_k$ and $q(x) = x$ or $q(x) = \ell_{j_k} - x$ (46) becomes

$$\begin{aligned} & -\frac{\ell_{j_k}}{2} |\partial_x^2 u_n^{j_k}(a_k)|^2 + \operatorname{Re} \left(d_{kj_k} \partial_x^2 u_n^{j_k}(a_s) \partial_x \overline{u_n^{j_k}}(a_s) \right) + \frac{1}{2} \int_0^{\ell_{j_k}} \beta_n^2 |u_n^{j_k}|^2 dx + \\ (47) \quad & \frac{3}{2} \int_0^{\ell_{j_k}} |\partial_x^2 u_n^{j_k}|^2 dx \rightarrow 0 \end{aligned}$$

where a_s is the end of e_{j_k} different from a_k .

Multiplying (42) by $\frac{1}{\beta_n^{1/2}} e^{-\beta_n^{1/2}(\ell_{j_k}-x)}$ or by $\frac{1}{\beta_n^{1/2}} e^{-\beta_n^{1/2}x}$, then, as in [25], after noting that $\partial_x^3 u_n^{j_k}(a_k)$ and $\beta_n u_n^{j_k}(a_k)$ tend to 0, we obtain

$$(48) \quad \partial_x^2 u_n^{j_k}(a_k) \longrightarrow 0.$$

Hence, (47) can be rewritten as

$$(49) \quad \operatorname{Re} \left(d_{kj_k} \partial_x^2 u_n^{j_k}(a_s) \partial_x \overline{u_n^{j_k}}(a_s) \right) + \frac{1}{2} \int_0^{\ell_{j_k}} \beta_n^2 |u_n^{j_k}|^2 dx + \frac{3}{2} \int_0^{\ell_{j_k}} |\partial_x^2 u_n^{j_k}|^2 dx \longrightarrow 0.$$

Now let $q \equiv 1$ (46) can be rewritten, for j_k , as

$$\begin{aligned} & -\frac{1}{2}\beta_n^2 |u_n^{j_k}(a_s)|^2 + \operatorname{Re} \left(\partial_x^3 u_n^{j_k}(a_s) \partial_x \overline{u_n^{j_k}}(a_s) \right) - \frac{1}{2} |\partial_x^2 u_n^{j_k}(a_s)|^2 - \\ & \operatorname{Re} \left(\mathbf{i}\beta_n f_n^{j_k}(a_s) \overline{u_n^{j_k}}(a_s) \right) \rightarrow 0. \end{aligned}$$

For j in $I(a_s)$, multiplying (42) by $\frac{1}{\beta_n^{1/2}} e^{-\beta_n^{1/2}x}$ or by $\frac{1}{\beta_n^{1/2}} e^{-\beta_n^{1/2}(\ell_j-x)}$, we get,

$$\frac{1}{\beta_n^{1/2}} d_{sj} \partial_x^3 u_n^j(a_s) + \varepsilon \partial_x^2 u_n^j(a_s) + \beta_n^{1/2} d_{sj} \partial_x u_n^j(a_s) + \varepsilon \beta_n u_n^j(a_s) \longrightarrow 0.$$

with $\varepsilon \in \{-1, 1\}$. Summing over $j \in I(a_s)$, by taking into account the continuity condition of u_n and $\partial_x^2 u_n$, the damping conditions, four in (6) and fifth in (7), and boundary condition at a_1 if $j = 1$, we deduce

$$\partial_x^2 u_n^j(a_s) + \beta_n u_n^j(a_s) \longrightarrow 0,$$

which leads to

$$\frac{1}{\beta_n^{1/2}} \partial_x^3 u_n^j(a_s) + \beta_n^{1/2} \partial_x u_n^j(a_s) = \alpha_n^j \longrightarrow 0.$$

Hence, for any positive real number a we have

$$\begin{aligned} \operatorname{Re} \left(\partial_x^3 u_n^j(a_s) \partial_x \overline{u_n^j(a_s)} \right) &= \operatorname{Re} \left(\frac{\partial_x^3 u_n^j(a_s)}{\beta_n^{1/2}} \beta_n^{1/2} \partial_x \overline{u_n^j(a_s)} \right) \\ &= \operatorname{Re} \left((\alpha_n^j - \beta_n^{1/2} \partial_x u_n^j(a_s)) \beta_n^{1/2} \partial_x \overline{u_n^j(a_s)} \right) \\ &\leq -\beta_n |\partial_x u_n^j(a_s)|^2 + \frac{a}{2} \beta_n |\partial_x u_n^j(a_s)|^2 + \frac{1}{2a} |\alpha_n^j|^2. \end{aligned}$$

Moreover, for any real positive number b we have

$$-\operatorname{Re} \left(i \beta_n f_n^j(a_s) \overline{u_n^j(a_s)} \right) \leq \frac{b}{2} \beta_n^2 |u_n^j(a_s)|^2 + \frac{1}{2b} |f_n^j(a_s)|^2.$$

Thus, we obtain the following framing for j_k ,

$$\begin{aligned} &-\frac{1}{2} \beta_n^2 |u_n^{j_k}(a_s)|^2 + \operatorname{Re} \left(\partial_x^3 u_n^{j_k}(a_s) \partial_x \overline{u_n^{j_k}(a_s)} \right) - \frac{1}{2} |\partial_x^2 u_n^{j_k}(a_s)|^2 \\ &\quad - \operatorname{Re} \left(i \beta_n f_n^{j_k}(a_s) \overline{u_n^{j_k}(a_s)} \right) - \frac{1}{2b} |f_n^{j_k}(a_s)|^2 \\ &\leq \left(-\frac{1}{2} + \frac{b}{2}\right) \beta_n^2 |u_n^{j_k}(a_s)|^2 - \frac{1}{2} |\partial_x^2 u_n^{j_k}(a_s)|^2 + \left(-1 + \frac{a}{2}\right) \beta_n |\partial_x u_n^{j_k}(a_s)|^2 \leq 0 \end{aligned}$$

with $a = b = \frac{1}{2}$. Which implies that

$$\beta_n^2 |u_n^{j_k}(a_s)|^2, |\partial_x^2 u_n^{j_k}(a_s)|^2, \beta_n |\partial_x u_n^{j_k}(a_s)|^2, \frac{1}{\beta_n} |\partial_x^3 u_n^{j_k}(a_s)|^2 \longrightarrow 0$$

and then all the expressions

$$\operatorname{Re} \left(\partial_x^3 u_n^{j_k}(a_s) \partial_x \overline{u_n^{j_k}(a_s)} \right), \operatorname{Re} \left(\partial_x^2 u_n^{j_k}(a_s) \partial_x \overline{u_n^{j_k}(a_s)} \right) \text{ and } \operatorname{Re} \left(i \beta_n f_n^{j_k}(a_s) \overline{u_n^{j_k}(a_s)} \right)$$

tend to 0 as n goes to infinity. Then (49) leads to

$$\frac{1}{2} \int_0^{\ell_{j_k}} \beta_n^2 |u_n^{j_k}|^2 dx + \frac{3}{2} \int_0^{\ell_{j_k}} |\partial_x^2 u_n^{j_k}|^2 dx \longrightarrow 0.$$

We iterate such procedure to obtain that for $j \neq 1$,

$$\frac{1}{2} \int_0^{\ell_j} \beta_n^2 |u_n^j|^2 dx + \frac{3}{2} \int_0^{\ell_j} |\partial_x^2 u_n^j|^2 dx \longrightarrow 0$$

and for $j \in I(a_2) - \{1\}$,

$$\beta_n^2 |u_n^j(a_2)|^2, |\partial_x^2 u_n^j(a_2)|^2, \operatorname{Re} \left(\partial_x^3 u_n^j(a_2) \partial_x \overline{u_n^j(a_2)} \right), \operatorname{Re} \left(\partial_x^2 u_n^j(a_2) \partial_x \overline{u_n^j(a_2)} \right) \rightarrow 0.$$

Finally let $j = 1$. Using continuity conditions of u_n and $\partial_x^2 u_n$, and damping conditions, four in (6) and fifth in (7), (46) leads to

$$\frac{1}{2} \int_0^{\ell_1} \beta_n^2 |u_n^1|^2 dx + \frac{3}{2} \int_0^{\ell_1} |\partial_x^2 u_n^1|^2 dx \longrightarrow 0.$$

In conclusion $\|y_n\|$ converge to 0, which contradicts the hypothesis that $\|y_n\| = 1$. $\square$

3.3. Polynomial stability. In this section we suppose that there is at least a beam following a string (Figure 2). We will prove that the solution of the whole system $(\mathcal{S})$ is polynomially stable and not exponentially stable.

Theorem 8. *If at least one beam follows a string from the root to a leaf (which is equivalent to say that the beam is located between two strings or a string and a leaf) then, the $\mathcal{C}_0$ -semigroup $(T(t))_{t \geq 0}$ is polynomially stable. More precisely: if all the sets of beams following strings are singletons then there is $C > 0$ such that*

$$\|e^{t\mathcal{A}}y_0\| \leq \frac{C}{t} \|y_0\|_{\mathcal{D}(\mathcal{A})}$$

for every $y_0 \in \mathcal{D}(\mathcal{A})$.

If there is a string followed by at least two beams then there is $C > 0$ such that

$$\|e^{t\mathcal{A}}y_0\| \leq \frac{C}{t^{2/3}} \|y_0\|_{\mathcal{D}(\mathcal{A})}$$

for every $y_0 \in \mathcal{D}(\mathcal{A})$.

Proof. In the sequel α is equal to 1 or $3/2$.

It suffices to prove that (17) holds. Suppose the conclusion is false. Then there exists a sequence (β_n) of real numbers, without loss of generality, with $\beta_n \rightarrow +\infty$, and a sequence of vectors $(y_n) = (u_n, v_n)$ in $\mathcal{D}(\mathcal{A})$ with $\|y_n\|_{\mathcal{H}} = 1$, such that

$$\|\beta_n^\alpha (\mathbf{i}\beta_n I - \mathcal{A})y_n\|_{\mathcal{H}} \rightarrow 0$$

which is equivalent to

$$(50) \quad \beta_n^\alpha (\mathbf{i}\beta_n u_n^j - v_n^j) = f_n^j \rightarrow 0, \quad \text{in } H^1(0, \ell_j), \quad j \text{ in } I_S,$$

$$(51) \quad \beta_n^\alpha (\mathbf{i}\beta_n u_n^j - v_n^j) = f_n^j \rightarrow 0, \quad \text{in } H^2(0, \ell_j), \quad j \text{ in } I_B,$$

$$(52) \quad \beta_n^\alpha (\mathbf{i}\beta_n v_n^j - \partial_x^2 u_n^j) = g_n^j \rightarrow 0, \quad \text{in } L^2(0, \ell_j), \quad j \text{ in } I_S,$$

$$(53) \quad \beta_n^\alpha (\mathbf{i}\beta_n v_n^j + \partial_x^4 u_n^j) = g_n^j \rightarrow 0, \quad \text{in } L^2(0, \ell_j), \quad j \text{ in } I_B.$$

Then

$$(54) \quad -\beta_n^\alpha (\beta_n^2 u_n^j + \partial_x^2 u_n^j) = g_n^j + \mathbf{i}\beta_n f_n^j, \quad j \text{ in } I_S,$$

$$(55) \quad \beta_n^\alpha (-\beta_n^2 u_n^j + \partial_x^4 u_n^j) = g_n^j + \mathbf{i}\beta_n f_n^j, \quad j \text{ in } I_B,$$

and

$$\|v_n^j\|^2 - \beta_n^2 \|u_n^j\|^2 \rightarrow 0, \quad j = 1, \dots, N.$$

Since $\operatorname{Re}(\langle \beta_n^\alpha (\mathbf{i}\beta_n - \mathcal{A})y_n, y_n \rangle_{\mathcal{H}}) = \sum_{a_k \in \mathcal{V}_{ext}^*} \beta_n^\alpha |v_n^{j_k}(a_k)|^2$, we obtain

$$\beta_n^{\frac{\alpha}{2}} |v_n^{j_k}(a_k)| \rightarrow 0, \quad \text{for } a_k \in \mathcal{V}_{ext}^*.$$

Then $\forall a_k \in \mathcal{V}_{ext}^B$, $\beta_n^{\frac{\alpha}{2}} \partial_x^3 u_n^{j_k}(a_k) \rightarrow 0$, $\forall a_k \in \mathcal{V}_{ext}^S$, $\beta_n^{\frac{\alpha}{2}} \partial_x u_n^{j_k}(a_k) \rightarrow 0$, $\forall a_k \in \mathcal{V}_{ext}^*$, $\beta_n^{1+\frac{\alpha}{2}} u_n^{j_k}(a_k) \rightarrow 0$, and recall that $\partial_x u_n^{j_k}(a_k) = 0$ for $a_k \in \mathcal{V}_{ext}^B$.

As in the proof of the previous theorem, if e_j is a string followed by no beam then

$$(56) \quad -\frac{1}{2}\beta_n^\alpha \int_0^{\ell_j} (|\partial_x u_n^j(x)|^2 + \beta_n^2 |u_n^j(x)|^2) dx \rightarrow 0,$$

and if a_k is an end of such edge then

$$(57) \quad \beta_n^{1+\frac{\alpha}{2}} u_n^j(a_k) \rightarrow 0 \quad \text{and} \quad \beta_n^{\frac{\alpha}{2}} \partial_x u_n^j(a_k) \rightarrow 0.$$

Now we can suppose without loss of generality that all edges are beams, except that related to the root which is a string.

Again, as in the previous proof, we have that for every beam e_j which not adjacent to the string,

$$(58) \quad \beta_n^{1+\frac{\alpha}{2}} \|u_n^j\|, \beta_n^{\frac{\alpha}{2}} \|\partial_x^2 u_n^j\| \longrightarrow 0,$$

and if a_k is an end of e_j then

$$(59) \quad \beta_n^{1+\frac{\alpha}{2}} |u_n^j(a_k)| \longrightarrow 0, \beta_n^{\frac{\alpha}{2}} |\partial_x^2 u_n^j(a_k)| \longrightarrow 0,$$

and all the expressions

$$Re \left(\beta_n^\alpha \partial_x^3 u_n^j(a_k) \partial_x \overline{u_n^j(a_k)} \right), Re \left(\beta_n^\alpha \partial_x^2 u_n^j(a_k) \partial_x \overline{u_n^j(a_k)} \right) \text{ and } Re \left(i \beta_n^{1+\alpha} f_n^j(a_k) \overline{u_n^j(a_k)} \right)$$

tend to zero as n goes to infinity.

Let $j \in I(a_2) - \{1\}$ (equivalently, e_j is a beam attached to a_2), then the real part of the inner product of (55) by $q \partial_x u_n^j$, with $q = x$ or $q = \ell_j - x$ gives

$$(60) \quad Re \left(\beta_n^\alpha d_{kj} \partial_x^2 u_n^j(a_2) \partial_x \overline{u_n^j(a_2)} \right) + \frac{1}{2} \beta_n^\alpha \int_0^{\ell_j} \beta_n^2 |u_n^j|^2 dx + \frac{3}{2} \beta_n^\alpha \int_0^{\ell_j} |\partial_x^2 u_n^j|^2 dx \longrightarrow 0$$

Summing over j in $I(a_2) - \{1\}$, one can deduce easily that

$$\sum_{j \in I(a_2) - \{1\}} \left(\frac{1}{2} \beta_n^\alpha \int_0^{\ell_j} \beta_n^2 |u_n^j|^2 dx + \frac{3}{2} \beta_n^\alpha \int_0^{\ell_j} |\partial_x^2 u_n^j|^2 dx \right) \longrightarrow 0$$

and in particular, for every beam e_j

$$\beta_n^\alpha \left(\int_0^{\ell_j} \beta_n^2 |u_n^j|^2 dx + \int_0^{\ell_j} |\partial_x^2 u_n^j|^2 dx \right) \longrightarrow 0$$

Now we want prove that, for j in $I(a_2) - \{1\}$, $\beta_n u_n^j(a_2)$ and $\partial_x^3 u_n^j(a_2)$ converge to 0.

First case: In this case there is one and only one beam e_j attached to e_1 and $\alpha = 1$.

The real part of the inner product of (55) by $q \partial_x u_n^j$, with $q = 1$ gives

$$(61) \quad -\frac{1}{2} \beta_n^3 |u_n^j(a_2)|^2 - \frac{1}{2} \beta_n |\partial_x^2 u_n^j(a_2)|^2 \rightarrow 0,$$

then $\beta_n^3 |u_n^j(a_2)|^2$ and $\beta_n |\partial_x^2 u_n^j(a_2)|^2$ converge to zero as n goes to infinite.

Multiplying (55) by $\frac{1}{\beta_n} e^{-\beta_n^{1/2} x}$ or by $\frac{1}{\beta_n} e^{-\beta_n^{1/2}(\ell_j - x)}$, we obtain, using the results below

$$\partial_x^3 u_n^j(a_2) \longrightarrow 0.$$

We deduce, using the fifth condition in (7), that $\partial_x u_n^1(a_1) \longrightarrow 0$.

Second case: In this case there is a string followed by at least two beams and $\alpha = 3/2$. First, (55) implies

$$(62) \quad \frac{\|\partial_x^4 u_n^j\|}{\beta_n^{1/4}} \longrightarrow 0.$$

Second, we need the following lemma (due to Gagliardo and Nirenberg [20]): $\square$

Lemma 9. (1) *There are two positive constants C_1 and C_2 such that for any w in $H^1(0, \ell_j)$,*

$$(63) \quad \|w\|_\infty \leq C_1 \|\partial_x w\|^{1/2} \|w\|^{1/2} + C_2 \|w\|.$$

(2) *There are two positive constants C_3 and C_4 such that for any w in $H^2(0, \ell_j)$,*

$$(64) \quad \|\partial_x w\| \leq C_3 \|\partial_x^2 w\|^{1/2} \|w\|^{1/2} + C_4 \|w\|.$$

The remainder of the proof. Applying the previous lemma several times, one can deduce easily that

$$(65) \quad \partial_x^3 u_n^j(a_1) \longrightarrow 0.$$

Return back to the fifth condition in (7) with using (65), we get $\partial_x u_n^1(a_1) \longrightarrow 0$. Now multiplying (54), when $u^j = u^1$, with $q(x) = \ell_1 - x$ or $q(x) = x$, then as for (43) and (44), with using (65) again and the continuity condition of u at a_2 , it follows

$$-\frac{1}{2} \int_0^{\ell_1} \left(|\partial_x u_n^1(x)|^2 + \beta_n^2 |u_n^1(x)|^2 \right) dx \longrightarrow 0.$$

Then we conclude that $\|y_n\| \rightarrow 0$ which contradicts the fact that $\|y_n\| = 1$ and the proof is then complete. $\square$

Now we consider a reduced system composed of one string e_1 and one beam e_2 such that $\ell_1 = \ell_2 = \pi$ and with control is applied on the beam. Precisely we consider the system

$$(\mathcal{S}_0) : \begin{cases} u_{tt}^1 - u_{xx}^1 = 0 \text{ in } (0, \pi) \times (0, \infty), \\ u_{tt}^2 + u_{xxxx}^2 = 0 \text{ in } (0, \pi) \times (0, \infty), \\ u^1(0, t) = u^2(0, t), \quad u_x^2(0, t) = 0, \quad u_{xxx}^2(0, t) = u_x^1(0, t), \\ u^1(\pi, t) = 0, \quad u_{xxx}^2(\pi, t) = u_t^2(\pi, t), \quad u_x^2(\pi, t) = 0, \\ u^j(x, 0) = u_0^j(x), \quad u_t^j(x, 0) = u_1^j(x), \quad j = 1, 2. \end{cases}$$

In view of Theorem 8 the system $(\mathcal{S}_0)$ is polynomial stable and we will prove that it is not exponentially stable. Note that if the control is applied on the string instead of the beam then the system is the exponential stability (by Theorem 7).

Theorem 10. *The system $(\mathcal{S}_0)$ is not exponential stable in the energy space $\mathcal{H}$.*

Proof. We prove that the corresponding semigroup $(T(t))_{t \geq 0}$ is not exponentially stable.

For $n \in \mathbb{N}$, such that $\sqrt{n}$ is integer and even let $\beta_n = n^2 + 2\sqrt{n} + \frac{1}{n}$ and $f_n = (0, 0, -\sin \beta_n x, 0)$, then $\beta_n \rightarrow +\infty$ and f_n is in $\mathcal{H}$ and is bounded. Let $y_n = (u_n^1, u_n^2, v_n^1, v_n^2) \in \mathcal{D}(\mathcal{A})$ such that $(\mathcal{A} - i\beta_n)y_n = f_n$. We will prove that $y_n \rightarrow +\infty$.

We have

$$(66) \quad \beta_n^2 u_n^1 + \partial_x^2 u_n^1 = \sin \beta_n x,$$

$$(67) \quad -\beta_n^2 u_n^2 + \partial_x^4 u_n^2 = 0.$$

then u_n^1 and u_n^2 are of the form

$$\begin{aligned} u_n^1 &= c_1 \sin(\beta_n x) + \left(-\frac{x}{2\beta_n} + c_2\right) \cos(\beta_n x), \\ u_n^2 &= d_1 \sin(\sqrt{\beta_n} x) + d_2 \cos(\sqrt{\beta_n} x) + d_3 \sinh(\sqrt{\beta_n} x) + d_4 \cosh(\sqrt{\beta_n} x). \end{aligned}$$

The transmission and boundary conditions are rewritten as follows

$$(68) \quad d_2 + d_4 = c_2,$$

$$(69) \quad \sqrt{\beta_n}(d_1 + d_3) = 0,$$

$$(70) \quad \beta_n^{3/2}(-d_1 + d_3) = -\frac{1}{2\beta_n} + \beta_n c_1,$$

and

$$(71) \quad c_1 \sin(\beta_n \pi) + \left(-\frac{\pi}{2\beta_n} + c_2\right) \cos(\beta_n \pi) = 0,$$

$$(72) \quad d_1 \cos(\sqrt{\beta_n} \pi) - d_2 \sin(\sqrt{\beta_n} \pi) + d_3 \cosh(\sqrt{\beta_n} \pi) + d_4 \sinh(\sqrt{\beta_n} \pi) = 0,$$

and

$$(73) \quad \begin{aligned} &\beta_n^{3/2}(-d_1 \cos(\sqrt{\beta_n} \pi) + d_2 \sin(\sqrt{\beta_n} \pi) + d_3 \cosh(\sqrt{\beta_n} \pi) + d_4 \sinh(\sqrt{\beta_n} \pi)) \\ &= i\beta_n(d_1 \sin(\sqrt{\beta_n} \pi) + d_2 \cos(\sqrt{\beta_n} \pi) + d_3 \sinh(\sqrt{\beta_n} \pi) + d_4 \cosh(\sqrt{\beta_n} \pi)). \end{aligned}$$

Summing (72) and (73), we obtain

$$(74) \quad \begin{aligned} &2\sqrt{\beta_n}(d_3 \cosh(\sqrt{\beta_n} \pi) + d_4 \sinh(\sqrt{\beta_n} \pi)) \\ &= i(d_1 \sin(\sqrt{\beta_n} \pi) + d_2 \cos(\sqrt{\beta_n} \pi) + d_3 \sinh(\sqrt{\beta_n} \pi) + d_4 \cosh(\sqrt{\beta_n} \pi)). \end{aligned}$$

Substituting (68) and (69) into (72) leads to

$$(75) \quad d_4 = \frac{\cosh(\sqrt{\beta_n} \pi) - \cos(\sqrt{\beta_n} \pi)}{\sinh(\sqrt{\beta_n} \pi) + \sin(\sqrt{\beta_n} \pi)} d_1 + \frac{\sin(\sqrt{\beta_n} \pi)}{\sinh(\sqrt{\beta_n} \pi) + \sin(\sqrt{\beta_n} \pi)} c_2.$$

Now, by substituting (68-70) and (75) into (74) we get

$$(76) \quad \begin{aligned} &2d_1 \left(\sqrt{\beta_n} h(\sqrt{\beta_n} \pi) - 2\beta_n \tan(\beta_n \pi) \sin(\sqrt{\beta_n} \pi) \sinh(\sqrt{\beta_n} \pi) \right. \\ &\quad \left. + i \left(1 - \cosh(\sqrt{\beta_n} \pi) \cos(\sqrt{\beta_n} \pi) + \sqrt{\beta_n} \tan(\beta_n \pi) h(\sqrt{\beta_n} \pi) \right) \right) \\ &= \left(2\sqrt{\beta_n} \sin(\sqrt{\beta_n} \pi) \sinh(\sqrt{\beta_n} \pi) - i h(\sqrt{\beta_n} \pi) \right) \left(-\frac{1}{2\beta_n^2} \tan(\beta_n \pi) + \frac{\pi}{2\beta_n} \right) \end{aligned}$$

with

$$h(\sqrt{\beta_n} \pi) = \cosh(\sqrt{\beta_n} \pi) \sin(\sqrt{\beta_n} \pi) + \sinh(\sqrt{\beta_n} \pi) \cos(\sqrt{\beta_n} \pi).$$

We want to prove that $\beta_n^{3/2} d_1$ is equivalent to $\sqrt{n} \frac{\pi^2}{2}$ as n goes to infinity.

Since $\beta_n = n^2 + 2\sqrt{n} + \frac{1}{n}$ then $\sqrt{\beta_n} = n + \frac{1}{\sqrt{n}} = n(1 + o(\frac{1}{\sqrt{n}}))$ and

$$\begin{aligned}\sin(\sqrt{\beta_n}\pi) &= \sin(\frac{\pi}{\sqrt{n}}) = \frac{\pi}{\sqrt{n}} + o(\frac{1}{\sqrt{n}}), \\ \cos(\sqrt{\beta_n}\pi) &= \cos(\frac{\pi}{\sqrt{n}}) = 1 + o(\frac{1}{\sqrt{n}}), \\ \tan(\beta_n\pi) &= \tan(\frac{\pi}{n}) = \frac{\pi}{n} + o(\frac{1}{n\sqrt{n}}), \\ \cosh(\sqrt{\beta_n}\pi) &= \cosh((n + \frac{1}{\sqrt{n}})\pi) = \frac{e^{n\pi}}{2}(1 + \frac{\pi}{\sqrt{n}} + o(\frac{1}{\sqrt{n}})), \\ \sinh(\sqrt{\beta_n}\pi) &= \frac{e^{n\pi}}{2}(1 + \frac{\pi}{\sqrt{n}} + o(\frac{1}{\sqrt{n}})),\end{aligned}$$

then

$$\begin{aligned}h(\sqrt{\beta_n}\pi) &= \frac{e^{n\pi}}{2}(1 + \frac{2\pi}{\sqrt{n}} + o(\frac{1}{\sqrt{n}})), \\ \sqrt{\beta_n} \sinh(\sqrt{\beta_n}\pi) \sin(\sqrt{\beta_n}\pi) \tan(\beta_n\pi) &= \frac{e^{n\pi}}{2}(\frac{\pi^2}{\sqrt{n}} + o(\frac{1}{\sqrt{n}})) \\ \sqrt{\beta_n} \sinh(\sqrt{\beta_n}\pi) \sin(\sqrt{\beta_n}\pi) &= \pi\sqrt{n}\frac{e^{n\pi}}{2}(1 + \frac{\pi}{\sqrt{n}} + o(\frac{1}{\sqrt{n}}))\end{aligned}$$

Hence, (76) implies

$$2d_1\sqrt{\beta_n}\frac{e^{n\pi}}{2}(1 + o(1)) = \frac{e^{n\pi}}{2}\sqrt{n}(2\pi + o(1))(-\frac{1}{2\beta_n^2}\tan(\beta_n\pi) + \frac{\pi}{2\beta_n}).$$

which leads to

$$(77) \quad 2\beta_n^{3/2}d_1 \sim_{+\infty} \pi^2\sqrt{n}.$$

Now suppose that $\partial_x u_n^1$ is bounded. The real part of the inner product of (66) with $(\pi - x)\partial_x u_n^1$ gives

$$-\frac{\pi}{2}\left|-\frac{1}{2\beta_n} + \beta_n c_1\right|^2 - \frac{\pi}{2}|\beta_n c_2|^2 = -\frac{1}{2}(\|u_n^1\|^2 + \|\partial_x u_n^1\|^2) + \operatorname{Re}\left(\int_0^\pi \sin(\beta_n x)(\pi - x)\partial_x \overline{u_n^1} dx\right).$$

By taking into account (69-70) and (77), $\beta_n^2 \|u_n^1\|^2 + \|\partial_x u_n^1\|^2$ converge to infinity. In conclusion y_n is not bounded. $\square$

Remark 11. Let $\varepsilon > 0$. By taking $\beta_n = n^2 + 2n^{1-\alpha} + \frac{1}{n^{2\alpha}}$ with $0 < \alpha < \varepsilon$ and such that $n^{1-\alpha}$ is integer and even and y_n is such that $f_n = (\beta_n^{\frac{1}{2}-\varepsilon}(\mathcal{A} - i\beta_n))y_n$, then we can prove that y_n is not bounded and then the polynomial stability of $(\mathcal{S})$ can't be better than $\frac{1}{t^2}$.

COMMENT

If we replace the boundary conditions by the followings

$$\begin{aligned}u^1(a_1, t) &= 0, \quad (1 - \delta)u_{xx}^1(a_1, t) = 0, \\ u_{xx}^{jk}(a_k, t) &= -u_t^{jk}(a_k, t), \quad a_k \in \mathcal{V}_{ext}^S, \\ u_{xx}^{jk}(a_k, t) &= -u_{tx}^{jk}(a_k, t), \quad u^{jk}(a_k, t) = 0, \quad a_k \in \mathcal{V}_{ext}^B,\end{aligned}$$

then we obtain the same results.

ACKNOWLEDGEMENTS

The authors thank the referees for many valuable remarks which helped us to improve the paper significantly.

REFERENCES

- [1] A. B. Abdallah and F. Shel, Exponential Stability of a General Network of 1- d Thermoelastic Materials, *Mathematical Control and Related Fields*, **2** (2012), 1-16.
- [2] K. Ammari, Asymptotic Behavior of some Elastic Planar Networks of Bernoulli-Euler Beams, *Applicable Analysis*, **86** (2007), 1529-1548.
- [3] K. Ammari and D. Jellouli and M. Mehrenberger, Feedback stabilization of a coupled string-beam system, *Netw. Heterog. media*, **4** (2009), 19-34.
- [4] K. Ammari and M. Jellouli, Stabilization of star-shaped networks of strings, *Differential and Integral Equations*, **17** (2004), 1395-1410.
- [5] K. Ammari and M. Jellouli and M. Khenissi, Stabilization of generic trees of strings, *J. Dynamical. Control. Systems*, **11** (2005), 177-193.
- [6] K. Ammari and M. Mehrenberger, Study of the nodal feedback stabilization of a string-beams network, *J Appl Math Comput.*, **36** (2012), 441-458.
- [7] K. Ammari, D. Mercier, V. Régnier and J. Valein, Spectral analysis and stabilization of a chain of serially Euler-Bernoulli beams and strings, *Commun. Pure Appl. Anal*, **11** (2012), 785-807.
- [8] K. Ammari and S. Nicaise, Stabilization of a transmission wave/plate equation, *J. Differential Equations*, **249** (2010), 707-727.
- [9] K. Ammari and S. Nicaise, *Stabilization of elastic systems by collocated feedback*, *Lectures Notes in Mathematics*, 2124, Springer, Cham, 2015.
- [10] C. Batty and T. Duyckaerts, Non-uniform stability for bounded semigroups on Banach spaces, *J. Evol. Equ.*, **8** (2008), 765-780.
- [11] J. Von Below, A characteristic equation associated to an eigenvalue problem on c^2 -networks, *Lin. Algebra Appl.*, **71** (1985), 309-325.
- [12] A. Borichev and Y. Tomilov, Optimal polynomial decay of functions and operator semigroups, *Math. Ann.*, **347** (2010), 455-478.
- [13] R. Dàger and E. Zuazua, *Wave propagation, observation and control in 1-d flexible multi-structures*, Springer-Verlag, Berlin, 2006, Mathématiques & Applications.
- [14] B. Dekoninck and S. Nicaise, The eigenvalue problem for networks of beams, *Linear Algebra and its Applications*, **314** (2000), 165-189.
- [15] L. M. Gearhart, Spectral theory for contraction semigroups on Hilbert space, *Trans. Amer. Math. Soc.*, **236** (1978), 385-394.
- [16] Z.J. Han and L. Wang, Riesz Basis Property and Stability of Planar Networks of Controlled Strings, *Acta Appl. Math.*, **110** (2010), 511-533.
- [17] F. L. Huang, Characteristic condition for exponential stability of linear dynamical systems in Hilbert spaces, *Ann. Diff. Eqs.*, **1** (1985), 43-56.
- [18] J.E. Lagnese, G. Leugering and E. J. P. G. Schmidt, *Modelling, Analysis and Control of Dynamic Elastic Multi-link Structure*, System & Control: Fundation & Applications, Birkhauser, Boston, MA, 1994.
- [19] Z. Liu and B. Rao, Characterization of polynomial decay rate for the solution of linear evolution equation, *Z. Angew. Math. Phys.*, **56** (2005), 630-644.
- [20] Z. Liu and S. Zheng, *Semigroups associated with dissipative systems*, Chapman & Hall/CRC, 1999.
- [21] D. Mercier and V. Régnier, Control of a Network of Euler-Bernoulli Beams, *J. Math. Anal. Appl.*, **342** (2008), 874-894.
- [22] A. Pazy, *Semigroup of Linear Operators and Applications to Partial Differential Equations*, Springer, 1983.
- [23] J. Prüss, On the spectrum of C_0 -semigroups, *Trans. Amer. Math. Soc.*, **284**, 847-857.
- [24] F. Shel, Exponential Stability of a Network of Elastic and Thermoelastic Materials, *Math. Meth. in Appl. Sci.*, **36** (2013), 869-879.
- [25] F. Shel, Exponential Stability of a Network of Beams, *J. Dynam. Control Syst.*, **21** (2015), 443-460.

- [26] J. Valein and E. Zuazua, Stabilization of the wave equation on 1-d networks, *SIAM J. Control Optim.*, **48** (2009), 2771-2797.
- [27] B. Z. Guo and J. M. Wang, Riesz Basis and Stabilization for the Flexible Structure of a Symmetric Tree-Shaped beam Network, *Math. Meth. in Appl. Sci.*, **31** (2008), 289-314.

UR ANALYSIS AND CONTROL OF PDE, UR 13ES64, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES OF MONASTIR, UNIVERSITY OF MONASTIR, 5019 MONASTIR, TUNISIA
E-mail address: `kais.ammari@fsm.rnu.tn`

UR ANALYSIS AND CONTROL OF PDE, UR 13ES64, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES OF MONASTIR, UNIVERSITY OF MONASTIR, 5019 MONASTIR, TUNISIA, TUNISIA
E-mail address: `farhat.shel@ipeit.rnu.tn`